

# Band ratio review – in a nutshell

Band ratioing = Divide the DNs in one band by the DNs in another band for each pixel to create a new data channel

Purposes:


- a. Spectral slope enhancement – highlight the difference between two bands
- b. Suppress / reduce topographic effect – shadow
- c. Include ratio as channel input for classifications
- d. Threshold for thematic analysis / display
- e. Generate useful colour composites (geology)

The NIR/ Red ratio highlights the rapid increase in NIR reflection by healthy veg. ->

Why is NIR/Red ‘better’ than NIR/Green ?  
Is it better than NIR / Blue ?

**Related image arithmetic:**

- Band addition, subtraction, multiplication



# Indices

**Ratios** ... enhance albedo contrasts by reducing inter-band similarities

e.g. Near-IR / Red ... to identify vegetation

---

Some sources refer to ratios as indices:

Ratio Vegetation Index (RVI) = **NIR / Red** > 1 = vegetated

\* RVI can create very high values (if Red Band DN is low)

Difference Vegetation Index (DVI) = **NIR-Red** > 0 = vegetated

\* DVI is heavily influenced by different lighting

‘Combining’ these two creates the most common vegetation index, involving subtraction, addition and division ... and ‘normalization’


# **Normalised** Difference Vegetation Index: NDVI

Proposed in the 1960s with Tiros/Nimbus weather satellites

$$NDVI = \frac{(NIR - RED)}{(NIR + RED)}$$

**Max= 1.0**

**Min= -1.0**




# Normalised Difference Vegetation Index NDVI

Division compensates for differential illumination

Normalisation yields values -1 to 1  
... stored in a 32 bit channel

= a close estimate of **biomass**  
also referred to as **greenness**

enhances the vegetation signal, while  
minimizing the impact of solar radiation and bare soil



- Negative values of NDVI (values approaching -1) correspond to water.
- Positive values close to zero (0 to 0.1) = barren areas of rock, sand, or snow.
- low, positive values represent shrub and grassland (0.2 to 0.5)
- high values indicate temperate and tropical rainforests (0.6 to 0.9)

What do you get when you store NDVI values in an 8-bit or 16-bit channel ?? Nothing!

# Highest NDVI values in different environments

<https://medium.com/regen-network/remote-sensing-indices-389153e3d947>


Ecosystem	Typical NDVI values	Location	References
Boreal forest	0.6-0.8	Alaska	Parent and Verbyla, 2010
Temperate forest	0.3-0.7	France	Pettorelli et al., 2006
Coastal rainforest	0.88-0.92	Solomon Islands	Garonna et al., 2009
Alpine pastures	0-0.35	Italy	Pettorelli et al., 2007
Annual grassland	0.15-0.45	California	Gamon et al., 1995
Desert	0.06-0.12	Sinai, Egypt	Dall'Olmo and Karnieli, 2002

**Table 1** - Typical NDVI values for different ecosystems (Pettorelli, 2013)

Landsat browser: <https://earthnow.usgs.gov/observer>




# Annual and interannual changes in NDVI



## Canada

NDVI values increase with 'green-up' in spring  
Peak in mid-late July ..  
start to drop in August,  
and into the fall

From mid-summer  
-> Near-IR decreases  
-> Red increases (till leaf fall)  
-> Implications for wildlife  
habitat and movement





# Special sensors for NDVI

SPOT 5 has extra bands / wide sensor in visible/NIR with 1 km resolution to capture a repeat 2400 km swath for global coverage

MODIS and NOAA-AVHRR have 250m/1000m red /near-IR bands for NDVI

NDVI is used to measure vegetation amount or biomass, in regional and global estimates. NDVI is directly related to photosynthesis and thus energy absorption of plant canopies





NDVI Anomaly



## Monitoring monthly and yearly changes, and anomalies in NDVI

Long term changes:  
may represent global impacts e.g. large scale forest change - clearance / regrowth

The difference between the average NDVI for a particular month of a given year (such as August 1993, above) and the average NDVI for the same month over the last 20 years is the NDVI anomaly. In 1993, heavy rain in the Northern Great Plains led to flooding in the Missouri River. The resulting lush vegetation appears as a positive anomaly (green). MODIS



Many satellite sensors specifically have red and Near IR bands to assess global vegetation

Table 1. Low or no-cost satellite sensors and data streams utilized for land surface phenology studies<sup>1</sup>

Sensor	Satellite	Overpass/ Orbit Frequency	Data Source (terrestrial data)	Data Record (years)	Spatial Resolution(s)	Processed Time Step	Latency
AVHRR	NOAA series	Daily	USGS/EROS <sup>2</sup>	1989- present	1 km	1-week, 2-week	~24 hours
AVHRR	NOAA series	Daily	NASA Ecocast <sup>1</sup>	1982-2013	8 km	Twice monthly	N/A
MSS	Landsat 1-5	18 days	USGS/EROS <sup>2</sup>	1972-1992	79 m	Distributed by scene	N/A
TM	Landsat 4-5	16 days	USGS/EROS <sup>2</sup>	1982-2011	30 m	Distributed by scene	N/A
ETM+	Landsat 7	16 days	USGS/EROS <sup>2</sup>	1999- present	30 m	Distributed by scene	~1-3 days
Vegetation	SPOT	1-2 days	VITO <sup>4</sup>	1999- present	1.15 km	10-day	~3 months
MODIS	Terra	1-2 days	LPDAAC <sup>5</sup>	2000- present	250 m, 500 m, 1 km	8-day, 16- day	~7-30 days
MODIS	Aqua	1-2 days	LPDAAC <sup>5</sup>	2002- present	250 m, 500 m, 1 km	8-day, 16- day	~7-30 days

# MODIS: MODerate-resolution Imaging Spectroradiometer

36 spectral bands ranging in wavelength 0.4  $\mu\text{m}$  to 14.4  $\mu\text{m}$  and at spatial resolutions 250m to 1km. Swath: 2330 km Earth covered 1-2 days

Designed to combine some of the characteristics of AVHRR and Landsat TM  
= 'Superspectral' (vs Multispectral)

MODIS is THE main medium resolution EO satellite sensor, but nearing end of life – gradually replaced by the Visible Infrared Imaging Radiometer Suite (VIIRS) onboard Suomi (Finland)


Reflected Solar Bands			Emissive Bands
<i>Aggregated 250 m</i>	<i>Aggregated 500 m</i>	<i>1 km</i>	<i>1 km</i>
Band 1 (620-670 nm)	Band 3 (459-479 nm)	Band 8 (405-420 nm)	Band 20 (3.660-3.840 $\mu\text{m}$ )
Band 2 (841-876 nm)	Band 4 (545-565 nm)	Band 9 (438-448 nm)	Band 21 (3.929-3.989 $\mu\text{m}$ )
	Band 5 (1230-1250 nm)	Band 10 (483-493 nm)	Band 22 (3.939-3.989 $\mu\text{m}$ )
	Band 6 (1628-1652 nm)	Band 11 (526-536 nm)	Band 23 (4.020-4.080 $\mu\text{m}$ )
	Band 7 (2105-2155 nm)	Band 12 (546-556 nm)	Band 24 (4.433-4.498 $\mu\text{m}$ )
		Band 13L (662-672 nm)	Band 25 (4.482-4.549 $\mu\text{m}$ )
		Band 13H (662-672 nm)	Band 27 (6.535-6.895 $\mu\text{m}$ )
		Band 14L (673-683 nm)	Band 28 (7.175-7.475 $\mu\text{m}$ )
		Band 14H (673-683 nm)	Band 29 (8.400-8.700 $\mu\text{m}$ )
		Band 15 (743-753 nm)	Band 30 (9.580-9.880 $\mu\text{m}$ )
		Band 16 (862-877 nm)	Band 31 (10.780-11.280 $\mu\text{m}$ )
		Band 17 (890-920 nm)	Band 32 (11.770-12.270 $\mu\text{m}$ )
		Band 18 (931-941 nm)	Band 33 (13.185-13.485 $\mu\text{m}$ )
		Band 19 (915-965 nm)	Band 34 (13.485-13.785 $\mu\text{m}$ )
		Band 26 (1.360-1.390 $\mu\text{m}$ )	Band 35 (13.785-14.085 $\mu\text{m}$ )
			Band 36 (14.085-14.385 $\mu\text{m}$ )

# MODIS (Moderate-resolution Imaging Spectroradiometer)

EarthExplorer datasets

## MODIS SPECIAL THEMES

MODIS Team Member	MODIS Product
E. Vermote	Surface Reflectance
Z. Wan	Land Surface Temperature
A. Strahler/J.-P. Muller	BRDF/Albedo
A. R. Huete/C. O. Justice	Vegetation Indexes
R. B. Myneni/S. W. Running	LAI/FPAR
C. O. Justice/Y. J. Kaufman	Fires/Burned Area
D. Hall	Snow/Ice/Sea Ice
J. R. G. Townshend/A. Strahler	Land Cover/Land Cover Change
S. W. Running	PSN/NPP



Launched by NASA in 1999 on the [Terra](#) (EOS AM) satellite, and in 2002 on the [Aqua](#) (EOS PM) satellite.


Terra: 10.30am descending

Aqua: 1.30pm ascending

# Delineation of Grizzly Bear Habitat in Bute Inlet

GEOG357 project

- Assumes bears are attracted to highest biomass areas  
e.g. avalanche slopes  
*Sieved NDVI threshold*





# Utilization of Landsat TM and Digital Elevation Data for the Delineation of Avalanche Slopes in Yoho National Park (Canada)

K. Wayne Forsythe and Roger D. Wheate






Fig. 1. Avalanche Slope at Takakkaw Falls, Yoho National Park. Looking west, slope is approximately 500 metres wide.

Avalanche slopes in black



**Example Vineyards**  
**Standard NDVI**

NDVI Vigor



'backwards'  
colour ramp !

Green GNDVI (NDGI) := (NIR-G) / (NIR+G)

**GNDVI** is more sensitive to chlorophyll variation in the crop than NDVI. It can be used in crops with dense canopies or in more advanced stages of development while NDVI is more suitable for estimating crop vigor during the early stages.







Soil-adjusted Vegetation Index (**SAVI**) =  $1.5 * (NIR - R) / (NIR + R + 0.5)$

Optimised Soil-adjusted Vegetation Index (**OSAVI**) =  $(NIR - R) / (NIR + R + 0.16)$

MSAVI: Modified Soil Adjusted Vegetation Index (MSAVI) minimizes the effect of bare soil on the Soil Adjusted Vegetation Index (SAVI)


$$MSAVI2 = \frac{(2 * NIR + 1 - 1\sqrt{(2 * NIR + 1) - 8 * (NIR - RED)})}{2}$$





**NDWI (Water):**  $(\text{Green} - \text{NIR}) / (\text{Green} + \text{NIR})$  1996  
- Mapping water - lakes, floods etc..  $> 0.2 = \text{water}$

Also **NDMI** =  $(\text{NIR} - \text{SWIR1}) / (\text{NIR} + \text{SWIR1})$   
**Moisture** content in leaves (also sometimes referred to as NDWI)  
Note: reversed order of bands compared to NDVI





NDWI of Canadian lakes. Acquired on 2020-08-05, Sentinel-hub

NDWI of Italy. Acquired on 2020-08-01



$$\text{NDMI} = (\text{NIR} - \text{SWIR}) / (\text{NIR} + \text{SWIR})$$



Moisture content in leaves (also sometimes referred to as NDWI)



# Snow: $NDSI = (Green - SWIR) / (Green + SWIR)$

NDSI has mostly been used for assessments of

1. Snow cover detection and mapping
2. Discriminate snow and clouds
3. Detect glacier ice in shadowed terrain
4. Detect frozen lakes
5. Glacier mapping





# Red Edge indices – Sentinel-2 (2015) (and Worldview 2009)

$NDRE = (NIR - RED\ EDGE) / (NIR + RED\ EDGE)$  (Normalized Difference Red Edge):



It is closely related to NDVI; it enables to identify vigor decreases earlier notably in agriculture





# Normalised (Difference) Burn Ratio (Index)

NDBR  $(\text{NIR} - \text{SWIR2}) / (\text{NIR} + \text{SWIR2})$  Landsat 8 OLI:  $\text{NBR} = (5 - 7) / (5 + 7)$



<https://www.usgs.gov/landsat-missions/landsat-normalized-burn-ratio>





# GEOG 357 project 2020

Kinbasket Lake  
August 2018

Sentinel 2 MSI  
Bands: SWIR2, NIR, Red

NDBR:  
 $(\text{NIR} - \text{SWIR2}) / (\text{NIR} + \text{SWIR2})$

Note: we've found that NBRI  
does not work as well in drier  
places e.g. southern BC, as there  
was limited vegetation beforehand




# Normalized Difference Built-up Index

$$\text{NDBI} = (\text{SWIR} - \text{NIR}) / (\text{SWIR} + \text{NIR})$$

uses the NIR and SWIR bands  
to emphasize built-up areas.  
(=lack of vegetation)

São José dos Campos, Brazil

**Figure 10.** NDBI calculated from the image of TM/Landsat 5 for 1986, 2001, and 2010.



[https://www.researchgate.net/publication/233629918\\_A\\_Quantitative\\_Approach\\_for\\_Analyzing\\_the\\_Relationship\\_between\\_Urban\\_Heat\\_Islands\\_and\\_Land\\_Cover](https://www.researchgate.net/publication/233629918_A_Quantitative_Approach_for_Analyzing_the_Relationship_between_Urban_Heat_Islands_and_Land_Cover)



## Summary of Indices

Vegetation and soil: NDVI most common

with many variants e.g. NDGI, DVI, SAVI, NRDE, NDBI

Water: NDWI

Snow: NDSI

Burns (fire): NDBR

All have values = -1.0 to +1.0 – stored in 32-bit real channel

This standardization enables direct comparison for change detection

# Many more examples... latest count > 150 ?

Intro+apps: [https://www.icimod.org/capacity-building/2020/totafg/day2/remote\\_sensing\\_indices\\_pt.pdf](https://www.icimod.org/capacity-building/2020/totafg/day2/remote_sensing_indices_pt.pdf)

Intro+6: <https://medium.com/regen-network/remote-sensing-indices-389153e3d947>

ENVI: <https://www.nv5geospatialsoftware.com/docs/AlphabeticalListSpectralIndices.html>

index database (300): <https://www.indexdatabase.de>

ArcGIS: <https://enterprise.arcgis.com/en/portal/10.7/use/band-arithmetic-function.htm>

ArcGIS: <https://pro.arcgis.com/en/pro-app/latest/help/data/imagery/indices-gallery.htm>

Sentinel: <https://custom-scripts.sentinel-hub.com/custom-scripts/sentinel-2/indexdb/>